

Kentucky Department for Medicaid Services

Pharmacy and Therapeutics Advisory Committee Recommendations

July 21, 2011 Meeting

The following chart provides a summary of the recommendations that were made by the Pharmacy and Therapeutics Advisory Committee at the July 21, 2011 meeting. Review of the recommendations by the Secretary of the Cabinet for Health and Family Services and final decisions are pending.

	Description of Recommendation	P & T Vote
1	<u>New Products to Market: vandetanib</u> Place this product preferred with similar quantity limits in the PDL class titled Oral Oncology Agents.	Passed 8 For 0 Against
2	<u>New Products to Market: Viibryd®</u> Place this product preferred in the PDL class titled Antidepressants: SSRIs; however, only approve Viibryd® after trial and failure of one SSRI.	Passed 8 For 0 Against
3	<u>New Products to Market: Zytiga™</u> Place this product preferred in the PDL class titled Oral Oncology Agents; however, only approve in combination with prednisone for a diagnosis of metastatic castration-resistant prostate cancer (CRPC) after: <ul style="list-style-type: none">• A trial of chemotherapy with docetaxel or mitoxantrone; OR• If the patient has a poor performance status.	Passed 8 For 0 Against
4	<u>New Products to Market: Horizant®</u> Horizant® should be approved for a diagnosis of restless legs syndrome (RLS) after trial and failure of ONE of the following: <ul style="list-style-type: none">○ Levodopa/carbidopa, OR○ Pramipexole, OR○ Ropinirole	Passed 6 For 2 Against

	Description of Recommendation	P & T Vote
5	<p><u>New Products to Market: Victrelis™</u></p> <p>Victrelis™ should be approved for a diagnosis of hepatitis C (CHC) genotype 1 infection after the patient has received 4 weeks of ribavirin and peginterferon therapy if they are receiving concurrent therapy with ribavirin and peginterferon. Victrelis™ should have a quantity limit of 12 capsules per day and be limited to one course of therapy per lifetime. Durations of therapy should be based on the following:</p> <p>a. Cirrhosis or previous treatment with peginterferon / ribavirin with documented lack of achievement of > 2 log reduction at week 12 with previous treatment:</p> <ol style="list-style-type: none"> Approve for 14 weeks After 14 weeks of therapy: <ol style="list-style-type: none"> If HCV-RNA level is ≤ 100 IU/mL at week 12 of therapy, approve for 12 more weeks If HCV-RNA results at week 24 of therapy are undetectable, approve for an additional 18 weeks (44 weeks total therapy) If HCV-RNA results at week 24 are detectable, discontinue all 3 therapies (Victrelis™ and peginterferon/ribavirin). <p>b. If none of above in a:</p> <ol style="list-style-type: none"> Approve for 14 weeks If HCV-RNA level is ≤ 100 IU/mL at week 12 of therapy, approve for 12 more weeks After 26 weeks, continuation of therapy should be approved based on the following: <ol style="list-style-type: none"> Treatment naïve patients: <ol style="list-style-type: none"> If HCV-RNA results at week 8 and 24 are both undetectable – 2 more weeks then discontinue all 3 therapies (Victrelis™ and peginterferon/ribavirin) – total duration of Victrelis™ therapy = 28 weeks If HCV-RNA results at week 8 are detectable and week 24 are undetectable – 10 more weeks – total duration of Victrelis™ therapy = 36 weeks If HCV-RNA results at week 24 are detectable, discontinue all 3 therapies (Victrelis™ and peginterferon/ ribavirin). Previously treated or relapsed patients: <ol style="list-style-type: none"> If HCV-RNA results at week 8 and 24 are both undetectable – 10 more weeks (then discontinue all 3) – total duration of Victrelis™ therapy = 36 weeks If HCV-RNA results at week 8 are detectable and week 24 results are undetectable 10 more weeks – total duration of Victrelis™ therapy = 36 weeks If HCV-RNA results at week 24 are detectable, discontinue all 3 therapies (Victrelis™ and peginterferon/ribavirin). 	<p>Passed</p> <p>8 For</p> <p>0 Against</p>

	Description of Recommendation	P & T Vote
6	<u>New Products to Market: Incivek™</u> Incivek™ should be approved for a diagnosis of hepatitis C (CHC) genotype 1 infection if the patient is receiving concurrent therapy with ribavirin and peginterferon. Incivek™ should have a quantity limit of 6 tablets per day for a total duration of 12 weeks and be limited to one course of therapy per lifetime.	Passed 8 For 0 Against
7	<u>New Products to Market: Sylatron™</u> Allow the use of Sylatron™ for a diagnosis of melanoma only.	Passed 8 For 0 Against
8	<u>New Products to Market: Tradjenta™</u> Place this product non preferred with similar quantity limits in the PDL class titled Diabetes: DPP-4 Inhibitors, unless cost parity to preferred DPP-4 Inhibitors.	Passed 8 For 0 Against
9	<u>New Products to Market: Daliresp™</u> Place this product preferred with similar quantity limits in the PDL class titled Anticholinergics, Inhaled; however, only approve after trial and failure of an inhaled anticholinergic or long-acting bronchodilator.	Passed 8 For 0 Against
10	<u>New Products to Market: Natroba™</u> Place this product non preferred in the PDL class titled Topical Antiparasitics.	Passed 8 For 0 Against
11	<u>5-ASA Derivatives, Rectal Preparations</u> 1. DMS to select preferred agent (s) based upon economic evaluation; however, at least one unique chemical entity should be preferred. Both suppositories and enemas should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the 5-ASA Derivatives, Topical Preparations class, require a PA until reviewed by the P&T Advisory Committee.	Passed 8 For 0 Against
12	<u>5-ASA Derivatives, Oral Preparations</u> 1. DMS to select preferred agent (s) based upon economic evaluation; however, at least two unique chemical entities, one of which should be oral mesalamine, should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the 5-ASA Derivatives, Oral Preparations class, require a PA until reviewed by the P&T Advisory Committee.	Passed 8 For 0 Against

	Description of Recommendation	P & T Vote
13	<p><u>Anti-Migraine: 5-HT1 Receptor Agonists</u></p> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however, at least one unique chemical entity should be preferred. Additionally, at least one non-oral dosage form should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. Agents in this class should have quantity limits based on the FDA-approved maximum dose and duration. 4. As part of quantity limit override criteria, patients should be on concurrent migraine prophylaxis medication (beta blocker, tricyclic antidepressant, calcium channel blocker, etc.) at a therapeutic dose. 5. For any new chemical entity in the Anti-Migraine: 5-HT1 Receptor Agonists class, require a PA until reviewed by the P&T Advisory Committee. 	<p>Passed</p> <p>8 For 0 Against</p>
14	<p><u>Hematopoietic Agents</u></p> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based upon economic evaluation. 2. All hematopoietic agents should require Prior Authorization. 3. For any agent not selected as preferred, DMS should allow continuation of therapy if there is a paid claim in the past 90 days. 4. For any new chemical entity in the Hematopoietic Agents class, require a PA until reviewed by the PTAC. 	<p>Passed</p> <p>8 For 0 Against</p>
15	<p><u>Hematopoietic Agents Clinical Criteria</u></p> <p>Erythropoiesis stimulating agents should be approved for recipients meeting one of the following criteria:</p> <ul style="list-style-type: none"> • The patient has a hemoglobin of less than 12 g/dL AND one of the following diagnoses: <ul style="list-style-type: none"> ○ Anemia associated with chronic renal failure OR anemia associated with kidney transplantation; OR ○ Treatment of chemotherapy induced anemia for non-myeloid malignancies; OR ○ Drug-induced anemia (examples, not all inclusive: Retrovir® or Combivir® or ribavirin); OR ○ Autologous blood donations by patients scheduled to undergo nonvascular surgery. 	<p>Passed</p> <p>8 For 0 Against</p>

	Description of Recommendation	P & T Vote
16	<u>Multiple Sclerosis Agents</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however, at least glatiramer, one interferon β-1b and one interferon β-1a product should be preferred. 2. Agents not selected as preferred will be considered non preferred and require PA. 3. Place quantity limits on these products based on maximum recommended dose. 4. Agents in this class should have no higher than a tier 2 copay regardless of PDL status. 5. For any new chemical entity in the Multiple Sclerosis Agents class, require a PA and quantity limit until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
17	<u>Ampyra™ Clinical Criteria</u> After 12 weeks of therapy (84 days), Ampyra™ therapy will be allowed to continue if the diagnosis is multiple sclerosis and Ampyra™ has shown clinical efficacy.	Passed 8 For 0 Against
18	<u>Oral Antiemetics: Anticholinergics</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however at least three unique chemical entities should be preferred. Promethazine and prochlorperazine should be among the preferred agents. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the Oral Anti-Emetics: Anticholinergics class, require a PA until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
19	<u>Oral Antiemetics: 5-HT₃ Antagonists</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however at least one unique chemical entity should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. Quantity limits should be removed from all oral dosages forms in this class. 4. For any new chemical entity in the Oral Anti-Emetics: 5-HT₃ Antagonists, require a PA until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
20	<u>Sancuso® Clinical Criteria</u> Sancuso® should be approved if the patient is currently undergoing cancer chemotherapy and one of the following is true: <ul style="list-style-type: none"> ○ The provider wishes to use this product to avoid the need for IV anti-emetics; OR ○ There has been a trial/failure on one preferred product. 	Passed 8 For 0 Against

	Description of Recommendation	P & T Vote
21	<u>Oral Antiemetics: NK-1 Antagonists</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however at least one unique chemical entity should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the Oral Anti-Emetics: NK₁ antagonist, require a PA until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
22	<u>Oral Antiemetics: Δ-9-THC Derivatives</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however at least one unique chemical entity should be preferred. 2. All agents in this category should require Prior Authorization to prevent miss-use. 3. For any new chemical entity in the Oral Anti-Emetics: Δ-9-THC Derivatives require a PA until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
23	<u>Oral Antiemetics: Δ-9-THC Clinical Criteria</u> Cannabinoids will be approved if one of the following is true: <ol style="list-style-type: none"> 1. Nausea and vomiting associated with cancer chemotherapy AFTER failure to respond adequately to at least ONE other anti-emetic therapy; OR 2. Anorexia associated with weight loss in patients with AIDS or cancer (dronabinol ONLY). 	Passed 8 For 0 Against
24	<u>H₂ Receptor Antagonists</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however, at least two unique chemical entities should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the H₂ Receptor Antagonists class, require a PA until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
25	<u>Anti-Ulcer Protectants</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based upon economic evaluation; however, at least two unique chemical entities should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the Anti-Ulcer Protectants class, require a PA until reviewed by the P&T Advisory Committee. 	Passed 8 For 0 Against
26	<u>Combination Products for H. pylori</u> <ol style="list-style-type: none"> 1. DMS to select preferred agent (s) based on economic evaluation; however, at least Prevpac® should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. Agents in this class should have quantity limits based on the FDA-approved maximum dose. 4. For any new chemical entity in the Combination Products for H. pylori class, require a PA until reviewed by the P&T Advisory Committee. 	Passed 5 For 3 Against

	Description of Recommendation	P & T Vote
27	<u>Antispasmodics / Anticholinergics</u> 1. DMS to select preferred agent (s) based on economic evaluation. However, at least one formulation of atropine, dicyclomine, glycopyrrolate, hyoscyamine, methscopolamine, and scopolamine should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the Antispasmodics / Anticholinergics class, require a PA until reviewed by the P&T Advisory Committee.	Passed 8 For 0 Against
28	<u>Antidiarrheals</u> 1. DMS to select preferred agent (s) based on economic evaluation; however at least two unique chemical entities should be preferred. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the Antidiarrheals class, require a PA until reviewed by the P&T Advisory Committee.	Passed 8 For 0 Against
29	<u>Laxatives and Cathartics</u> 1. DMS to select preferred agent (s) based on economic evaluation; however, at least four unique chemical entities should be preferred. The preferred products should include lactulose, polyethylene glycol, and one agent used for bowel evacuation or colon cleansing. 2. Agents not selected as preferred will be considered non-preferred and will require Prior Authorization. 3. For any new chemical entity in the Laxatives and Cathartics class, require a PA until reviewed by the P&T Advisory Committee.	Passed 8 For 0 Against
30	<u>Amitiza[®] Clinical Criteria</u> Amitiza [®] should be approved for the following diagnoses: <ul style="list-style-type: none"> • Irritable Bowel Syndrome with constipation; OR • Chronic Idiopathic Constipation after failure of one laxative. 	Passed 8 For 0 Against
31	<u>Relistor[®] Clinical Criteria</u> Relistor [®] should be approved if all of the following criteria are met: <ul style="list-style-type: none"> • Diagnosis of opioid-induced constipation; AND • Patient has advanced illness, which is defined as a terminal disease (incurable cancer or other end-stage disease); AND • Trial and failure (unless contraindicated or intolerant to) of an agent in each of the following drug classes: <ul style="list-style-type: none"> ○ Stool softening agent; AND ○ Peristalsis-inducing agent. 	Passed 8 For 0 Against